



Isle of Wight
NHS Trust

Patient Information Leaflet

When your baby dies before birth

Produced by:
Maternity Department

Date: June 2017
Review due: June 2020



**This page is left
intentionally
blank**

If you are unable to read this leaflet because English is not your first language, please ask someone who speaks English to telephone PALS on 01983 534850 for further information and help.

Polish:

Jeśli nie jesteś w stanie przeczytać tej ulotki bo angielski nie jest pierwszym językiem, poproś kogoś, kto mówi po angielsku, o kontakt telefoniczny z (PALS 01983 534850) aby uzyskać więcej informacji i pomoc

Russian:

Если вы не можете прочитать этот буклет на английском языке потому что не является первым языке, пожалуйста, попросите кого-нибудь, кто говорит на английском языке для телефонного (PALS 01983 534850) для получения дополнительной информации и помощи

Turkish:

Eğer İngilizce ana diliniz değilse, çünkü bu broşürü okumak için yapamıyorsanız, daha fazla bilgi için 01983 534850 üzerinde PALS telefon İngilizce bilen birine sormak ve yardım lütfen

Bulgarian:

Ако не сте в състояние да прочетете тази листовка, тъй като английският не е първи език, опитайте някой, който говори английски, за да телефонирам (PALS на 01983 534850) за повече информация и помощ

Czech:

Pokud nejste schopni přečíst tuto příbalovou informaci, protože angličtina není vaším rodným jazykem, zeptejte se někoho, kdo mluví anglicky na telefonní PALS na 01983 534850 pro další informace a pomoc

Bengali:

আপনি ইংরেজি আপনার প্রথম ভাষা না থাকার কারণে এই লফিলটে পড়তে অক্ষম হন, তাহলে আরও তথ্যের জন্য 01983 534850 নভেগিশেন PALS টলেফি এন ইংরেজি কথা কটে জিজ্ঞাসা করুন এবং সাহায্য করুন

Introduction

Being told that your baby has died before birth is devastating. The grief experienced by parents, and by close relatives and friends, is also extremely distressing. You and your family will be given considerable support by your midwives and doctors; however, it can be hard to take in all the information given to you at this time and therefore it is often helpful to have it in writing as well.

This leaflet helps to explain the care you will receive during and after the birth of your baby, and care in future pregnancies. It does not cover everything related to the longer-term emotional support you may need. Sands, the stillbirth and neonatal death charity, offers support to anyone affected by the death of a baby before, during or shortly after birth. If at any time you feel uncertain about anything, the healthcare professionals looking after you will be there to help.

Why does it happen?

The question that every parent wishes to have answered is 'why did my baby die?'. It is not always possible to give an answer but a cause is found more than two-thirds of the time. A baby dying before birth occurs in one in every 200 pregnancies.

The most common reason for a baby dying in the womb is because the baby has not been growing properly. There are other causes, including infection, abnormal development of the baby, diabetes, early separation of the placenta and pre-eclampsia (high blood pressure and protein in your urine). You and your partner will be offered tests to try to find out why your baby died (see below).

How is the death of your baby before labour confirmed?

Death of a baby in the womb is confirmed by an ultrasound scan. The scan is able to show if your baby's heart has stopped beating. If you wish, you can ask for another scan to reconfirm your baby's death.

Sometimes, after it has been confirmed that your baby has died, you may still feel as if your baby is moving. This is caused by the shifting movements of your baby within the fluid in your womb and can be very distressing. Your midwife will be able to talk to you about this.

What happens next?

Experienced doctors and midwives will talk to you and your partner about the death of your baby. They will explain your choices for birth, and the various tests that may be offered. You and your partner will be given support and plenty of time to make any decisions you need to make.

You will be seen by a doctor to make sure that you are well yourself. This includes, for example, checking for infection and pre-eclampsia.

What are my choices for birth?

Your doctor will discuss with you and your partner the different options of when and how to have your baby. The advice will depend on your general health, your pregnancy and any previous birth experiences, and also your personal wishes. You will be looked after by an experienced midwife and, where possible, you will be cared for in an individual room that has facilities for your partner or another companion to stay with you.

Inducing labour (starting labour with medication)

You may choose to have labour induced as soon as possible after confirmation of your baby's death, or you may prefer to go home for a short while before treatment. Some women find it comforting to spend some time at home. There are different ways of inducing labour and your doctor will discuss with you which method would be the most appropriate for you. Often, a tablet is given by mouth 48 hours before admission for induction.

Letting nature take its course – You may choose to wait at home for labour to start naturally. If your waters have not broken and you are well physically, you are unlikely to come to any harm if you delay labour for a short period of time (up to 48 hours). Most women will go into labour naturally within 3 weeks of their baby dying in the womb.

You need to be aware that delaying the onset of labour will affect the appearance of your baby at birth. In addition, tests that you agree to being carried out on your baby may give less information. If you decide to wait for more than 48 hours, you will be advised to have a check-up at the hospital twice a week as a small number of women will become unwell during this time.

Your midwife or doctor will give you a contact number to ring if you have any queries or concerns while waiting for labour to start. You must ring your maternity unit if you have pain, bleeding or a smelly vaginal discharge, or if you feel unwell in any way.

You may be advised against delaying labour if you have pre-eclampsia or an infection, or if some of the blood tests show that a delay would put your health at risk.

You may decide that it would cause you greater anxiety to delay starting labour than having your labour induced (when labour is started with medication).

How will I give birth to my baby?

Vaginal birth is usually the recommended way to give birth to your baby. Although you may find the thought of a vaginal birth distressing, you may want to consider that:

- there are fewer risks to you;
- you will be able to go home more quickly;
- your recovery is likely to be quicker and more straightforward future pregnancies are less likely to have complications.

Your doctor and midwife will discuss your choices with you and your wishes will be respected. All types of pain relief will usually be available in labour. An epidural may not be possible if you have an infection or problems with blood clotting but all steps will be taken to ensure that you get the pain relief you need.

What if I have had a caesarean section before?

A consultant obstetrician should discuss your choices with you. If you have had one previous caesarean section, it is usually safe to have labour induced although it is not completely without risk. A vaginal birth not only has the advantages stated above (see How will I give birth to my baby?), but having a repeat caesarean section has additional risks and makes a future vaginal delivery less likely.

What happens when my baby is born?

You will be given additional support by the midwives and doctors looking after you, who know that giving birth is going to be a distressing experience for you and your family. You and your partner (and sometimes other members of your family, with your permission) may wish to see and/or hold your baby immediately after birth. You may like to wait to see your baby until a little later after birth. You may decide not to see your baby at all.

If you wish to see and hold your baby, then you will have the opportunity to spend as much time as you wish with your baby. A cold cot is also available to keep your baby's body cool and allows you and your family to spend a longer period of time with your baby following birth.

Some parents decide to take their baby home from the hospital for a short time. This can give you the opportunity to spend time with your baby quietly and privately. You will also be able to take the cold cot home with you. You may be able to take your baby straight home with you but if a post-mortem is required you will not be able to take your baby home until after this. You can arrange to bring your baby home after the post-mortem by contacting the bereavement support midwives.

Mementoes such as photographs, hand and footprints and locks of hair, if possible, can be taken, and these are often valued by parents. A memory box will also be offered to enable you to take home if you wish. If you do not wish to take these home with you straight away or if you are not sure about whether you would like to keep them, they can usually be kept securely in your hospital records (check with your midwife whether this is possible at your hospital).

You can then have them at a later date if you wish. It will be your choice as to what you would like to do.

What happens after my baby has been born?

A member of staff will talk to you and your partner about the funeral choices for your baby, and about registering the birth. Your religious and cultural considerations will be taken into account.

Grief for you, your partner and family following the death of a baby can be severe and last a long time. Options for support and bereavement counselling should be offered to you and your partner. Your GP will be informed and you and your partner will be given information on Sands and local support groups where available.

You can usually go home when you wish, although if you have been unwell you may be advised to stay in the maternity unit for a little longer. Some women and partners want to go home as soon as possible, whereas others prefer to stay for a while. You can talk to your midwife about the best time for you to go home.

After the birth of your baby you may experience the sensation of breast milk coming into your breasts, making them feel full and uncomfortable. Ice packs, breast support and pain relief may help but, despite this, one in three women may still experience excessive discomfort. In this instance, there are tablets available that will stop your body producing milk and relieve your symptoms at this distressing time. The hospital doctor or your GP can prescribe these tablets if you need them.

As with any birth, you will have bleeding and pain for a few days. This should settle down within a week and you will be checked over by your midwife at home. Problems are uncommon after birth but you should contact your GP or hospital if:

- **your bleeding gets heavier;**
- **you have pain in your abdomen that doesn't settle;**
- **you have a smelly vaginal discharge;**
- **you feel unwell or shivery;**
- **you have pain, swelling or soreness in your legs;**
- **you have shortness of breath or chest pain, or cough up blood.**

What tests / investigations will I be offered?

You will be offered tests for you and your baby that may help to find out why your baby has died. A cause is found in two-thirds of all stillbirths and this can help with planning care in a future pregnancy. Unfortunately, despite tests, sometimes deaths cannot be explained.

You and your partner will be given time to think about which tests you would like to have done and you will be supported in any choices you make. You will need to sign a consent form for some of the tests. You will be offered an appointment at a later date to discuss the results of any tests that you have consented to being done.

Tests you and your partner will be offered include:

- Tests to look for conditions in your pregnancy, for example pre-eclampsia, infection or any problems with your liver or kidneys.

- Blood tests to see if there is any underlying medical cause, for example diabetes or thyroid problems.
- Blood tests to see if there is an underlying condition that makes your blood more likely to clot (thrombophilia or antiphospholipid syndrome). Pregnancy can sometimes affect the results of these tests and therefore they may need to be done again 6 weeks later.
- Swabs taken from your vagina, cervix and placenta and from your baby to look for any signs of infection.
- A test of your baby's chromosomes that will involve taking a blood sample or skin/muscle sample from your baby. This test is important because approximately 6% (approximately one in 17) of stillborn babies have abnormal chromosomes. If this is the case you and your partner may also be tested. You will be given further detailed written information about this test and you will need to give your consent and sign a form.
- A post-mortem examination for your baby and placenta, which can be limited or detailed as you wish. A post-mortem examination can provide very important information on why your baby has died. You will be given further detailed information about this and a chance to discuss your choices with a doctor; it will be your decision for your baby to have a post-mortem examination and you will need to give your consent and sign before this test can be done. Your individual rights and cultural and religious beliefs will always be respected.

- A detailed examination of your placenta even without a post-mortem examination may also provide valuable information.

What happens when I leave the Hospital?

You will receive a telephone call from a midwife the day after you return home. This is to make sure you are well physically and to see if you would like to have a home visit and/or if there is anything else that you need.

You will also be sent an appointment to see a Consultant Obstetrician. This appointment is an opportunity to discuss all the events surrounding your baby's death and to review the results of the tests that were performed prior to and after your delivery. Typically, this appointment is arranged for 6 weeks after you leave the hospital but can be up to 12 weeks depending on the tests and investigations that have been performed.

What extra care will I get if I become pregnant again?

If you decide to have another baby, you will usually be under consultant care and will be seen early in your pregnancy. You will usually have extra antenatal visits as you will understandably be anxious. You will be given additional support by the doctors and midwives looking after you throughout your pregnancy, who will be aware of your previous loss. The precise pattern of your care will be influenced by the results of any tests and whether a cause has been found for the death of your baby.

Further information and support

If you require any further support or information please do not hesitate to contact the midwives at the Maternity Unit by calling 534349/534392.

Sands is a stillbirth and neonatal death charity that offers emotional support and practical help to anyone affected by the death of a baby before, during or shortly after birth. As well as supporting mothers and partners, Sands can help other members of the family, especially grandparents and other children, as well as friends. Information and support can be found at:

Helpline: 020 7436 5881

Website: www.uk-sands.org

Address: Sands, 3rd Floor, 28 Portland Place, London W1B 1LY

**This page is left
intentionally
blank**

**This page is left
intentionally
blank**

If you have any queries or concerns about the content of this leaflet please call (01983) 524081 and ask to speak to the relevant department.

If you require this leaflet in another language, large print or another format, please contact the PALS Team, telephone **01983 534850**, who will advise you.

Valuables should not be brought into the hospital. If patients have to bring in valuable items they should ask a nurse to store them safely and request a receipt for the items. You may not be able to have the valuable items returned if the time of discharge from hospital is out of hours.

We are sorry but the Trust cannot accept responsibility for loss or damage to items not given for safe keeping.

You can get further information on all sorts of health issues online at: **www.nhs.uk**

For Health advice and out of hours GP service please call the NHS 111 service on: **111**

We Value Your Views On Our Service

If you wish to comment on the care which you, your relative or friend has received, we will be pleased to hear from you. Please speak to the person in charge of the ward, clinic or service in the first instance or ask them to contact the PALS Team. If you wish to contact them directly, telephone on **01983 534850**.

Alternatively you may prefer to write to:

Chief Executive
Isle Of Wight NHS Trust
St Mary's Hospital
Newport, Isle of Wight, PO30 5TG

You can also share any concerns you have about our services with the Care Quality Commission (CQC) on **03000 61 61 61** or at **enquiries@cqc.org.uk**

All NHS sites are no smoking areas.

If you would like help and advice to stop smoking please call: Freephone 0800 169 0 169 to talk to the NHS Smoking Helpline.